

Guidelines for Prescribing Psychotropic Medication to Preschool Age Children (3-5 Years Old)

Diagnosis	1 st Line Treatment	2 nd Line Treatment	3 rd Line Treatment
<p>ADHD</p> <p>Diagnostic Assessment /Screening Tool</p> <ul style="list-style-type: none"> ▪ ADHD Rating Scale – IV Preschool or Connors Early Childhood - EC 	<p>Psychotherapeutic Trial</p> <ul style="list-style-type: none"> ▪ Parent Behavior Training (PBT) interventions¹ 	<p>Methylphenidate/Dexmethylphenidate</p> <ul style="list-style-type: none"> ▪ Medication can be discontinued quickly ▪ Review family/child history of heart condition* <p><u>Side Effects</u>¹</p> <ul style="list-style-type: none"> ▪ Loss of appetite - severely underweight and children who have lost a significant amount of weight should be carefully monitored by a pediatrician for increased nutritional needs. ▪ Increased blood pressure and heart rate should be monitored closely by a pediatrician. ▪ Stomach and/or head ache ▪ Irritability/moodiness² ▪ Insomnia/sedation 	<p>Amphetamine Formulations</p> <ul style="list-style-type: none"> ▪ Medication can be discontinued quickly ▪ Review family/child history of heart condition* ▪ As effective as methylphenidate in older children but no good studies have been done in children under 5. <p><u>Side Effects</u>¹</p> <ul style="list-style-type: none"> ▪ The side effect profile is significantly greater in this age group than for Methylphenidate/Dexmethylphenidate³ ▪ Loss of appetite - severely underweight and children who have lost a significant amount of weight should be carefully monitored by a pediatrician for increased nutritional needs. ▪ Increased blood pressure and heart rate should be monitored closely by a pediatrician. ▪ Stomach and/or head ache ▪ Irritability/moodiness² ▪ Insomnia/sedation
4th Line Treatment			
<p>Alpha-Agonists</p> <ul style="list-style-type: none"> ▪ Careful consideration of age and body weight, initial low liquid doses ▪ If discontinuation is planned, these medications must be decreased slowly in increments. ▪ A higher dosing range may be needed if there are other significant diagnoses¹ ▪ Parent education about safe administration and monitoring - Never give more than prescribed, if dose missed do not double dose, do not discontinue this medication abruptly. <p><u>Side Effects</u></p> <ul style="list-style-type: none"> ▪ Sedation/REM suppression ▪ Irritability ▪ Headache ▪ Slow heart rate - bradycardia, hypotension ▪ Low blood pressure – monitor blood pressure and heart rate*** ▪ Drying effect (e.g. dry mouth, changes in vision, constipation) 		<p>Atomoxetine</p> <p><u>Side Effects</u>⁴</p> <ul style="list-style-type: none"> ▪ Mood Swings ▪ Sleepiness ▪ Decreased appetite ▪ Abdominal Pain ▪ Vomiting ▪ Headache 	

* If there is a family history of structural heart disease or an arrhythmia, or if the patient has a heart condition, the patient should have a baseline ECG. Contact the child’s PCP to discuss safety issues. For more complicated cardiac pathology, an echocardiogram or a cardiology consultation may be indicated.

** If the patient loses weight such that his/her weight drops 2 percentile lines on a standard growth curve or if his/her weight falls below the 3rd percentile, the medication should be discontinued. The child may need a referral for a growth delay evaluation.

*** A baseline ECG is not indicated unless the patient has a pre-existing arrhythmia or cardiac disease.

Guidelines for Prescribing Psychotropic Medication to Preschool Age Children (3-5 Years Old)

Diagnosis	1 st Line Treatment	2 nd Line Treatment	3 rd Line Treatment
<p>Anxiety</p> <p>Diagnostic Assessment /Screening Tool</p> <ul style="list-style-type: none"> ▪ Spence Preschool Anxiety Scale: Parent Report - free tool to help assess children ages 3-6 with anxiety. <p>Anxiety</p> <p>http://www.scasweb site.com/docs/scas-preschool-scale.pdf</p> <ul style="list-style-type: none"> ▪ Ages and Stages Questionnaire: Social Emotional (ASQ-SE) 	<p>Psychotherapeutic Trial</p> <ul style="list-style-type: none"> ▪ Behavioral therapy or preschool CBT⁵ for a minimum of 12 weeks ▪ Parenting intervention for anxiety without mood disorder ⁶ 	<p>Fluoxetine</p> <ul style="list-style-type: none"> ▪ Last resort intervention due to the high incidence of SSRI related side effects, specifically behavioral activation in young children - for severe symptoms⁷ ▪ Planned discontinuation after 6-9 months ▪ Given the sensitivity to side effects in the young children, increase dose slowly. <p><u>Side Effects</u></p> <ul style="list-style-type: none"> ▪ Headache ▪ Stomach ache ▪ Insomnia or increased motor activity ▪ Increased energy /unrestrained behavior may increase in younger children and children who also have ADHD or brain disorders ⁸ ▪ FDA - Black box warning: SSRIs increase the risk for suicidal thinking ▪ With use of Fluoxetine, please review interactions with any other medications the child is taking i.e. asthma medications, antibiotics, seizure medications etc. ▪ Decreased appetite and weight loss ▪ Sleep disturbance 	<p>Sertraline</p> <ul style="list-style-type: none"> ▪ Last resort intervention due to the high incidence of SSRI related side effects, specifically behavioral activation in young children - for severe symptoms⁷ ▪ Planned discontinuation after 6-9 months <p><u>Side Effects</u></p> <ul style="list-style-type: none"> ▪ Headache ▪ Stomach ache ▪ Insomnia or increased motor activity ▪ Increased energy /unrestrained behavior may increase in younger children and children who also have ADHD or brain disorders ⁸ ▪ FDA - Black box warning: SSRIs increase the risk for suicidal thinking ▪ With use of Fluoxetine, please review interactions with any other medications the child is taking i.e. asthma medications, antibiotics, seizure medications etc.

Guidelines for Prescribing Psychotropic Medication to Preschool Age Children (3-5 Years Old)

Diagnosis	1 st Line Treatment	2 nd Line Treatment	3 rd Line Treatment
<p>Autism Spectrum Disorder</p> <p>Diagnostic Assessment /Screening Tool</p> <ul style="list-style-type: none"> Child Autism Rating Scale Modified Checklist for Autism in Toddlers Revised (M-CHAT-R) https://www.m-chat.org/mchat.php 	<p>Psychotherapeutic Trial</p> <ul style="list-style-type: none"> Parent psychoeducation Early intervention to address ¹ <ul style="list-style-type: none"> Language Social development Adaptive functioning Reduction in repetitive behaviors Aggression Tantrums Self injury Hyperactivity Anxiety and Mood <p>Dysregulation (if significant comorbid problems, please refer to those disorders in this guideline)⁹ <ul style="list-style-type: none"> Sensory sensitivity ⁹ <ul style="list-style-type: none"> Behavioral Therapy ⁹ <p>Applied Behavioral Analysis (ABA) gold standard</p> </p>	<p>Irritability and Aggression</p> <p>Risperidone</p> <ul style="list-style-type: none"> FDA indication for irritability and aggression in children aged 5 to 16 years with autistic disorder and symptoms of aggression, self-injury, temper tantrums and mood swings ⁹ Should only be given to young children with severe symptoms of aggression, self-injury, temper tantrums and mood swings because: <ul style="list-style-type: none"> More severe symptoms showed greater improvement with Risperidone.¹⁰ The weight related/metabolic side effects of antipsychotic medication/risperidone ¹¹ <ul style="list-style-type: none"> Dietary education should be offered Metabolic monitoring is needed <p><u>Side Effects</u></p> <ul style="list-style-type: none"> Changes in fasting blood sugar cholesterol, blood pressure and abdominal fat (metabolic syndrome) Tremor, stiffness, changes in eye movement, drooling (extrapyramidal side effects) Increased levels of the hormone prolactin Extreme restlessness Close monitoring of patients is essential⁹ 	<p>Irritability and Aggression</p> <p>Aripiprazole*</p> <ul style="list-style-type: none"> Good results in school aged population but no preschool data Good treatment effects and comparatively mild side-effects to other atypical antipsychotics ¹² <p><u>Side Effects</u></p> <ul style="list-style-type: none"> Sedation Weight gain Changes in fasting blood sugar cholesterol, blood pressure and abdominal fat (metabolic syndrome) Severe restlessness Tremor, stiffness, changes in eye movement, drooling (extrapyramidal side effects) <p>Guanfacine/Clonidine</p> <p><u>Side Effects</u> ⁹</p> <ul style="list-style-type: none"> FDA indication for 6-17 years Good results in school aged population Sedation Weight gain Light headed and unsteady ⁷
		<p>Hyperactivity</p> <p>Methylphenidate</p> <ul style="list-style-type: none"> Medication can be discontinued quickly Review family/child history of heart condition* ASD children are more sensitive to medication, particularly stimulants, than children with only ADHD ⁹ <p><u>Side Effects</u></p> <ul style="list-style-type: none"> Loss of appetite - severely underweight and children who have lost a significant amount of weight should be carefully monitored by a pediatrician for increased nutritional needs. Increased blood pressure and heart rate should be monitored closely by a pediatrician. Stomach and/or head ache Irritability/moodiness ² Insomnia/sedation 	<p>Hyperactivity</p> <p>Alpha-Agonists</p> <ul style="list-style-type: none"> Careful consideration of age and body weight, initial low liquid doses If discontinuation is planned, these medications must be decreased slowly in increments. A higher dosing range may be needed if there are other significant diagnoses³ Parent education about safe administration and monitoring - Never give more than prescribed, if dose missed do not double dose, do not discontinue this medication abruptly. <p><u>Side Effects</u></p> <ul style="list-style-type: none"> Sedation Irritability Headache

Guidelines for Prescribing Psychotropic Medication to Preschool Age Children (3-5 Years Old)

Diagnosis	1 st Line Treatment	2 nd Line Treatment	3 rd Line Treatment
Autism Spectrum Disorder		Hyperactivity (continued)	Hyperactivity (continued)
		<ul style="list-style-type: none"> ▪ Agitation ▪ Abnormal movements⁹ such as vocal tics like constant throat clearing , coughing, making noises, etc. 	<ul style="list-style-type: none"> ▪ Slow heart rate ▪ Low blood pressure ¹³ – monitor blood pressure and heart rate*** Drying effect (e.g. dry mouth, changes in vision, constipation)
			4 th Line Treatment
			Hyperactivity Atomoxetine <u>Side Effects</u> <ul style="list-style-type: none"> ▪ Mood Swings ▪ Decreased appetite ▪ Sleepiness ▪ Abdominal Pain
Diagnosis	1 st Line Treatment	2 nd Line Treatment	3 rd Line Treatment
Autism Spectrum Disorder		Repetitive Behaviors	Repetitive Behaviors
		<u>Fluoxetine</u> <ul style="list-style-type: none"> ▪ Last resort intervention for severe symptoms ▪ Planned discontinuation after 6-12 months ▪ Given the sensitivity to side effects in the young children, increase dose slowly.¹⁴ <u>Side Effects</u> <ul style="list-style-type: none"> ▪ Headache ▪ Stomach ache ▪ Insomnia or increased motor activity ▪ Increased energy /unrestrained behavior may increase in younger children and children who also have ADHD or brain disorders ⁸ FDA - Black box warning: SSRIs increase the risk for suicidal thinking ▪ With use of Fluoxetine, please review interactions with any other medications the child is taking i.e. asthma medications, antibiotics, seizure medications etc. ▪ Decreased appetite and weight loss ▪ Sleep disturbance 	<u>Fluvoxamine, Citalapram, Sertraline and Escitalopram</u> <ul style="list-style-type: none"> ▪ Studies support use in children 6 years and above but there is no data supporting use in children under six ¹⁴

Guidelines for Prescribing Psychotropic Medication to Preschool Age Children (3-5 Years Old)

Diagnosis	1 st Line Treatment	2 nd Line Treatment	3 rd Line Treatment
<p>Bipolar</p> <p>Diagnostic Assessment /Screening Tool</p> <ul style="list-style-type: none"> Young Mania Rating Scales <p>Note: Bipolar is very difficult to diagnose in a preschool population because mood changes and extreme emotions are common in this age group. While a diagnosis can be made in children as young as 3 years old, Bipolar Disorder remains extremely rare in this population.¹⁵ Once diagnosed, address mania first, higher incidence of rapid (daily) cycling and mixed mania^{16,17}</p>	<p>Psychotherapeutic Trial</p> <ul style="list-style-type: none"> Parent Child Interaction Therapy (PCIT)⁶ 	<p><u>Risperidone</u></p> <ul style="list-style-type: none"> Due to the weight related/metabolic side effects of antipsychotic medications such as risperidone¹¹ <ul style="list-style-type: none"> Dietary education should be offered with a prescription Metabolic monitoring is needed <p><u>Side Effects</u></p> <ul style="list-style-type: none"> Sedation/fatigue Weight gain Changes in fasting blood sugar cholesterol, blood pressure and abdominal fat (metabolic syndrome) Headache Tremor, stiffness, changes in eye movement, drooling (extrapyramidal side effects) Increased levels of the hormone prolactin Severe restlessness Close monitoring of patients is essential¹⁶ 	<p><u>Aripiprazole</u>¹²</p> <ul style="list-style-type: none"> Good results in school aged population but no preschool data Good treatment effects and mild side-effects relative to other atypical antipsychotics¹² <p><u>Side Effects</u></p> <ul style="list-style-type: none"> Sedation Weight gain/increased appetite Changes in fasting blood sugar cholesterol, blood pressure and abdominal fat (metabolic syndrome) Severe restlessness GI disturbance Headache Tremor, stiffness, changes in eye movement, drooling (extrapyramidal side effects)^{16,18} <p>Quetiapine</p> <p><u>Side Effects</u></p> <ul style="list-style-type: none"> Sedation/decreased energy Weight gain Changes in fasting blood sugar cholesterol, blood pressure and abdominal fat (metabolic syndrome) Severe restlessness GI disturbance Headache Tremor, stiffness, changes in eye movement, drooling (extrapyramidal side effects)^{16,19}

Guidelines for Prescribing Psychotropic Medication to Preschool Age Children (3-5 Years Old)

Diagnosis	1 st Line Treatment	2 nd Line Treatment	3 rd Line Treatment
<p>Depression</p> <p>Diagnostic Assessment /Screening Tool</p> <ul style="list-style-type: none"> Preschool Feelings Checklist ²⁰ 	<p>Psychotherapeutic Trial</p> <ul style="list-style-type: none"> Psychotherapeutic Treatment modalities that address the parent-child relationship such as Parent Child Interaction Therapy-Emotion Development (PCIT-ED) ²¹ 	<p><u>Fluoxetine</u></p> <ul style="list-style-type: none"> Last resort intervention due to the high incidence of SSRI related side effects, specifically behavioral activation in young children - for severe symptoms ^{1,7,22} Planned discontinuation after 9 months at therapeutic dose Given the sensitivity to side effects in young children, increase dose slowly. <p><u>Side Effects</u></p> <ul style="list-style-type: none"> Headache Stomach ache Insomnia or increased motor activity Increased energy /unrestrained behavior may increase in younger children and children who also have ADHD or neurodevelopmental disorders ⁸ FDA - Black box warning: SSRIs increase the risk for suicidal thinking With use of Fluoxetine, please review interactions with any other medications the child is taking i.e. asthma medication, antibiotic, seizure medication etc. Decreased appetite and weight loss Sleep disturbance 	<p><u>Citalopram/Escitalopram</u></p> <ul style="list-style-type: none"> Last resort intervention due to the high incidence of SSRI related side effects, specifically behavioral activation in young children ^{7,23,24} Clinical experience suggests other SSRIs such as Citalopram and Escitalopram may be easier for preschool children to tolerate. However, with Citalopram can change the electrical conduction through the heart.
<p>Disruptive Behavior Disorder (DBD) and Aggression</p> <p>Diagnostic Assessment /Screening Tool</p> <p><i>Note: Treat the co-morbid disorders contributing to disruptive behavior first</i></p> <ul style="list-style-type: none"> Eyberg Child Behavior Inventory (ECBI) 	<p>Psychotherapeutic Trial</p> <ul style="list-style-type: none"> Preschool CBT Parent Child Interaction Therapy (PCIT), Incredible Years Program, Collaborative Problem Solving etc. ²⁵ Infant/Toddler Parent Programs i.e. Child Parent Interactive Therapy Classroom-Based Interventions Token Reward Systems 	<ul style="list-style-type: none"> Presence of Disruptive/Aggressive Behavior and any other major mental illness – treat other disorder first. If other major mental illness adequately treated and that medication and therapy is addressing Disruptive/Aggressive Behavior, continue treatment of other MI. If Disruptive/Aggressive Behavior Alone: <p><u>Risperidone</u> - Close monitoring of patients is essential</p> <ul style="list-style-type: none"> Antipsychotics are often used to augment psychotherapy. For <u>severe</u> aggression in preschool age children, an atypical antipsychotic can be prescribed ⁸ <p><u>Side Effects</u></p> <p>Changes in fasting blood sugar cholesterol, blood pressure and abdominal fat (metabolic syndrome)</p> <ul style="list-style-type: none"> Tremor, stiffness, changes in eye movement, drooling (extrapyramidal side effects) Increased levels of the hormone prolactin Extreme restlessness 	

Guidelines for Prescribing Psychotropic Medication to Preschool Age Children (3-5 Years Old)

Diagnosis	1 st Line Treatment	2 nd Line Treatment	3 rd Line Treatment
<p>Obsessive Compulsive Disorder (OCD)</p> <p>Diagnostic Assessment /Screening Tool</p> <ul style="list-style-type: none"> Spence Preschool Anxiety Scale: Parent Report - free tool to help assess children ages 3-6 with anxiety. <p>http://www.scasweb site.com/docs/scas-preschool-scale.pdf</p> <p>²⁶</p>	<p>Psychotherapeutic Trial</p> <ul style="list-style-type: none"> Cognitive Behavioral Therapy CBT using exposure and response prevention techniques and involving parents is recommended ²⁶ 	<p><u>Fluoxetine, Sertraline and Fluvoxamine</u></p> <ul style="list-style-type: none"> Last resort intervention due to the high incidence of SSRI related side effects, specifically behavioral activation in young children - for severe symptoms ^{15,7,23} Has been approved by the Food and Drug Administration (FDA) for the treatment of OCD in children. Fluoxetine is 8 years and above. Sertraline is 6 years and above. Fluvoxamine is 8 years and above. Using dose equivalents due to insufficient research for children ages 3-5. Insufficient evidence to recommend one medication over the other Extreme caution should be used with these medication for severe OCD in this age group. ²⁷ Given the sensitivity to side effects in the young children, increase dose slowly. Planned discontinuation after 6-8 months at therapeutic dose ²⁷ <p><u>Side Effects</u></p> <ul style="list-style-type: none"> Headache Stomach ache Insomnia or increased motor activity Increased energy /unrestrained behavior may increase in younger children and children who also have ADHD or brain disorders ⁸ FDA - Black box warning: SSRIs increase the risk for suicidal thinking With use of Fluoxetine and Sertraline, please review interactions with any other medications the child is taking i.e. asthma medications, antibiotics, seizure medications etc. Decreased appetite and weight loss Sleep disturbance 	

Guidelines for Prescribing Psychotropic Medication to Preschool Age Children (3-5 Years Old)

Diagnosis	1 st Line Treatment	2 nd Line Treatment	3 rd Line Treatment
PTSD	Psychotherapeutic Trial <ul style="list-style-type: none"> ▪ Child-parent psychotherapy (CPP) for a 6 month trial ¹ or preschool CBT for minimum of 12 weeks ²⁸ 	Psychopharmacological interventions are not recommended for children under 6 years based on a lack of research evidence. Talk to a DCFS Psychopharmacology program consultant if symptoms are severe and therapeutic interventions are ineffective.	
Sleep Disturbance Diagnostic Assessment Screening Tool <ul style="list-style-type: none"> ▪ Sleep Log 	Parent Education <ul style="list-style-type: none"> ▪ Home environment evaluation ▪ Sleep hygiene ▪ Restless leg syndrome ▪ Sleep Apnea ▪ Sleep problem associated with other mental health diagnoses ▪ Behavior Intervention (2-4 weeks) 	<u>Melatonin</u> ²⁹ <ul style="list-style-type: none"> ▪ May be appropriate when sleep disturbance is impacting well-being and daytime functioning of child and/or caregiver ▪ Over-the-counter ▪ Short term use, 1 month maximum before reassessment 	<u>Alpha-Agonists</u> <ul style="list-style-type: none"> ▪ Careful consideration of age and body weight, initial low liquid doses ▪ If discontinuation is planned, these medications must be decreased slowly in increments. ▪ Short term use, 1 month maximum before reassessment ▪ Parent education about safe administration and monitoring - Never give more than prescribed, if dose missed do not double dose, do not discontinue this medication abruptly. <u>Side Effects</u> ²⁹ <ul style="list-style-type: none"> ▪ Respiratory depression ▪ Decreased REM (deep sleep) ▪ Irritability ▪ Headache ▪ Slow heart rate ▪ Low blood pressure – monitor blood pressure and heart rate*** ▪ Slow heart rate - bradycardia, hypotention ▪ Drying effect (e.g. dry mouth, changes in vision, constipation)

*** A baseline ECG is not indicated unless the patient has a pre-existing arrhythmia or cardiac disease.

1. Gleason MM, Egger HL, Emslie GJ, et al. Psychopharmacological Treatment for Very Young Children: Contexts and Guidelines. *J Am Acad Child Adolesc Psychiatry*. 2007;46(12):1532-1572. doi:10.1097/chi.0b013e3181570d9e.
2. Charach A, Carson P, Fox S, Ali MU, Beckett J, Lim CG. Interventions for Preschool Children at High Risk for ADHD: A Comparative Effectiveness Review. *PEDIATRICS*. 2013;131(5):e1584-e1604. doi:10.1542/peds.2012-0974.
3. Luby JL. Handbook of Preschool Mental Health, Second Edition : Development, Disorders, and Treatment. <http://public.eblib.com/choice/publicfullrecord.aspx?p=4659183>. Published 2016.
4. Kratochvil CJ, Vaughan BS, Mayfield-Jorgensen ML, et al. A Pilot Study of Atomoxetine in Young Children with Attention-Deficit/Hyperactivity Disorder. *J Child Adolesc Psychopharmacol*. 2007;17(2):175-186. doi:10.1089/cap.2006.0143.
5. Geller DA, March J. Practice parameter for the assessment and treatment of children and adolescents with obsessive-compulsive disorder. *Focus*. 2012;10(3):360–373.
6. Luby JL. Treatment of anxiety and depression in the preschool period. *J Am Acad Child Adolesc Psychiatry*. 2013;52(4):346–358.
7. Zuckerman ML, Vaughan BL, Whitney J, et al. Tolerability of Selective Serotonin Reuptake Inhibitors in Thirty-Nine Children Under Age Seven: A Retrospective Chart Review. *J Child Adolesc Psychopharmacol*. 2007;17(2):165-174. doi:10.1089/cap.2007.0086.
8. Sakolsky D, Birmaher B. Pediatric anxiety disorders: management in primary care: *Curr Opin Pediatr*. 2008;20(5):538-543. doi:10.1097/MOP.0b013e32830fe3fa.
9. Kaplan G, McCracken JT. Psychopharmacology of autism spectrum disorders. *Pediatr Clin North Am*. 2012;59(1):175-187, xii. doi:10.1016/j.pcl.2011.10.005.
10. Arnold LE, Farmer C, Kraemer HC, et al. Moderators, mediators, and other predictors of risperidone response in children with autistic disorder and irritability. *J Child Adolesc Psychopharmacol*. 2010;20(2):83–93.
11. Scahill L, Jeon S, Boorin SJ, et al. Weight Gain and Metabolic Consequences of Risperidone in Young Children With Autism Spectrum Disorder. *J Am Acad Child Adolesc Psychiatry*. 2016;55(5):415–423.
12. Oh J, Chang JG, Lee SB, Song D-H, Cheon K-A. Comparison of Aripiprazole and Other Atypical Antipsychotics for Pediatric Bipolar Disorder: A Retrospective Chart Review of Efficacy and Tolerability. *Clin Psychopharmacol Neurosci*. 2013;11(2):72-79. doi:10.9758/cpn.2013.11.2.72.
13. Scahill L, Aman MG, McDougle CJ, et al. A Prospective Open Trial of Guanfacine in Children with Pervasive Developmental Disorders. *J Child Adolesc Psychopharmacol*. 2006;16(5):589–598.
14. West L, Brunssen SH, Waldrop J. Review of the evidence for treatment of children with autism with selective serotonin reuptake inhibitors. *J Spec Pediatr Nurs*. 2009;14(3):183–191.

15. Luby JL, Tandon M, Belden A. Preschool Bipolar Disorder. *Child Adolesc Psychiatr Clin N Am.* 2009;18(2):391-403. doi:10.1016/j.chc.2008.11.007.
16. Peruzzolo TL, Tramontina S, Rohde LA, Zeni CP. Pharmacotherapy of bipolar disorder in children and adolescents: an update. *Rev Bras Psiquiatr.* 2013;35(4):393-405. doi:10.1590/1516-4446-2012-0999.
17. Geller B, Tillman R, Bolhofner K. Proposed Definitions of Bipolar I Disorder Episodes and Daily Rapid Cycling Phenomena in Preschoolers, School-Aged Children, Adolescents, and Adults. *J Child Adolesc Psychopharmacol.* 2007;17(2):217-222. doi:10.1089/cap.2007.0017.
18. Findling RL, McNamara NK, Youngstrom EA, et al. An Open-Label Study of Aripiprazole in Children with a Bipolar Disorder. *J Child Adolesc Psychopharmacol.* 2011;21(4):345-351. doi:10.1089/cap.2010.0102.
19. Joshi G, Petty C, Wozniak J, et al. A prospective open-label trial of quetiapine monotherapy in preschool and school age children with bipolar spectrum disorder. *J Affect Disord.* 2012;136(3):1143-1153. doi:10.1016/j.jad.2011.09.042.
20. Luby JL, Heffelfinger AK, Mrakotsky C, Hessler MJ, Brown KM, Hildebrand T. Preschool major depressive disorder: preliminary validation for developmentally modified DSM-IV criteria. *J Am Acad Child Adolesc Psychiatry.* 2002;41(8):928–937.
21. Lenze SN, Pautsch J, Luby J. Parent-child interaction therapy emotion development: a novel treatment for depression in preschool children. *Depress Anxiety.* 2011;28(2):153-159. doi:10.1002/da.20770.
22. Ercan ES, Kandulu R, Akyol Ardic U. Preschool children with obsessive–compulsive disorder and fluoxetine treatment. *Eur Child Adolesc Psychiatry.* 2012;21(3):169-172. doi:10.1007/s00787-012-0244-2.
23. Hetrick SE, McKenzie JE, Cox GR, Simmons MB, Merry SN. Newer generation antidepressants for depressive disorders in children and adolescents. In: The Cochrane Collaboration, ed. *Cochrane Database of Systematic Reviews.* Chichester, UK: John Wiley & Sons, Ltd; 2012. <http://doi.wiley.com/10.1002/14651858.CD004851.pub3>. Accessed September 16, 2016.
24. Safer DJ, Zito JM. Treatment-emergent adverse events from selective serotonin reuptake inhibitors by age group: children versus adolescents. *J Child Adolesc Psychopharmacol.* 2006;16(1-2):159–169.
25. Luby J, Mrakotsky C, Stalets MM, et al. Risperidone in preschool children with autistic spectrum disorders: an investigation of safety and efficacy. *J Child Adolesc Psychopharmacol.* 2006;16(5):575–587.
26. Whiteside SPH, Gryczkowski MR, Biggs BK, Fagen R, Owusu D. Validation of the Spence Children’s Anxiety Scale’s obsessive compulsive subscale in a clinical and community sample. *J Anxiety Disord.* 2012;26(1):111-116. doi:10.1016/j.janxdis.2011.10.002.
27. Coskun M, Zoroglu S. Efficacy and safety of fluoxetine in preschool children with obsessive-compulsive disorder. *J Child Adolesc Psychopharmacol.* 2009;19(3):297–300.

28. Cohen JA. Treating acute posttraumatic reactions in children and adolescents. *Biol Psychiatry*. 2003;53(9):827-833. doi:10.1016/S0006-3223(02)01868-1.
29. Pelayo R, Yuen K. Pediatric Sleep Pharmacology. *Child Adolesc Psychiatr Clin N Am*. 2012;21(4):861-883. doi:10.1016/j.chc.2012.08.001.